

Understanding mouse bile acid formation: Is it time to unwind why mice and rats make unique bile acids?¹

Mats Rudling²

Metabolism Unit and KI/AZ Integrated Cardio Metabolic Center, Department of Medicine, Center for Innovative Medicine, Department of Biosciences and Nutrition, Karolinska Institute at Karolinska University Hospital Huddinge, S-141 86 Stockholm, Sweden

Current knowledge on bile acid metabolism is largely based on extrapolations from animal experiments where the mouse has taken a front position, greatly due to the development of techniques making it feasible to construct mouse models where specific functions are deficient or overexpressed. However, there are several major differences between mice and humans as regards bile acid metabolism that are important to recognize when interpreting data obtained from experiments on mice and extrapolating that data to humans. One such difference is that in mice as well as in rats, bile acids are hydroxylated at the 6- β position to form MuriCholic Acids (MCAs), a set of bile acids that is unique to these species. Although the long pathway for the synthesis of bile acids in humans has progressively been uncovered (1), the structure(s) responsible for the formation of MCAs has remained a dark area, although it has previously been concluded that chenodeoxycholic acid (CDCA) should be at least one substrate for the formation of MCAs (2). In the present issue of the *Journal of Lipid Research*, Takahashi et al. (3) report that the *Cyp2c70* gene is key for the synthesis of MCAs. By screening liver extracts from different knock-out mouse strains for the absence of MCAs, they found that *Cyp2c* gene cluster knock-out (*Cyp2c*-null) mice are completely deficient in MCAs. Of the 15 functional candidate genes within that gene cluster, the authors came to the conclusion that the *Cyp2c70* gene is necessary for the formation of α -MCA from CDCA as well as for the formation of β -MCA from ursodeoxycholic acid (UDCA). In liver extracts from WT mice, the formation of α -MCA from CDCA occurred with an ~40-fold higher affinity than the formation of β -MCA from UDCA. However, in line with previous findings on gallbladder bile (4) or liver extracts (5, 6), they found that in liver extracts from WT mice, β -MCA was 5-fold more abundant than α -MCA, and that the level of conjugated tauro (T) β -MCA was 20-fold higher than that of T α -MCA. It was proposed that, presumably, epimerization of 7 α -MCA into 7 β -MCA, possibly mediated by intestinal microbiota, might explain the higher abundance of β -MCA. The ultimate model to prove these results, knock-out mice for *Cyp2c70* solely, is now highly warranted.

This is an open access article under the [CC BY](#) license.

DOI 10.1194/jlr.C072876

ARE MCAS OF INTEREST IN HUMAN MEDICINE?

The long and complex pathways for the synthesis of bile acids are classic examples of where it has been recognized that the level of synthesis is suppressed by the end products made, the bile acids. How this occurred was not well understood until the discovery that the farnesoid X receptor (FXR) serves as a specific receptor for bile acids in this negative feedback regulation (7, 8). This knowledge has in turn led to an increased awareness that the FXR agonistic activity varies greatly between different bile acids. Thus, CDCA and deoxycholic acid (DCA) are clearly more potent than cholic acid (CA) and UDCA, the latter two being poor FXR activators when studied alone in vitro (8, 9). However, when UDCA is used in the presence of potent FXR activating agonists such as DCA or CDCA, it can dampen the FXR stimulation from the agonists (9). In this respect, the MCAs are of particular interest. Although never mentioned in the report by Makishima et al. (7), it could be seen that the CYP7A1 protein in HepG2 cells was dose-dependently increased when cells were exposed to MCAs. Later studies on germ-free mice have highlighted that such animals have an enlarged pool of bile acids rich in MCAs and that T β -MCA and T α -MCA can serve as antagonists to CDCA as determined with a coactivator recruitment assay with recombinant human FXR (10). Germ-free mice have been shown to have an improved resistance to high-fat feeding (11). Interestingly, cholic acid free *CYP8B1*^{-/-} mice and mice treated with antibiotics share several features with germ-free mice. They have an induced bile acid synthesis and increased expression of the intestinal ASBT protein, an enlarged pool of bile acids enriched in MCAs, and their intestinal absorption of dietary fat including cholesterol is reduced (12–14). *CYP8B1*^{-/-} mice are also less prone to develop steatosis (12). Further, feeding WT mice with α - or β -MCAs reduces the intestinal absorption of cholesterol from 38 to 11% (15).

The present identification of *Cyp2c70* as key for the synthesis of MCAs from CDCA or UDCA should now make it

¹See referenced article, *J. Lipid Res.* 2016, 57: 2130–2137.

²To whom correspondence should be addressed.
e-mail: mats.rudling@ki.se


feasible to synthesize MCAs in vivo in species other than rats and mice. One may thus speculate whether the production of significant quantities of MCAs by human intestinal microbiota or in liver may result in beneficial metabolic effects such as reduced body weight, improved insulin sensitivity, and reduced liver lipids. One important substrate for this, CDCA, is certainly highly available in humans.

Other conditions where it may be of interest to generate MCAs in humans are different situations with cholestasis associated with high systemic levels of the potentially toxic CDCA. In bile duct-ligated rats, the urinary excretion of the water soluble MCAs increases about 400-fold (16) to become the major excretory route for bile acids. In addition, bile acid synthesis is induced.

Although such approaches may at first appear attractive, the fact that administered bile acids are often rapidly converted in vivo may result in unexpected responses. Thus, the administration of CDCA to mice induces the formation of MCAs from CDCA that dampens the expected suppression of bile acid synthesis by this treatment, while on the other hand, administration of the weak FXR agonist CA suppresses CYP7A1 strongly, presumably due to the pronounced reduction of FXR antagonistic MCAs that is seen during such treatment (5, 6).

FURTHER UNDERSTANDING OF THE PHYSIOLOGIC FUNCTION OF MCAS IN RATS AND MICE

It is well known that mice and rats are relatively resistant to high-fat feeding and that mice with boosted levels of MCAs, such as germ-free, *Cyp8b1*^{-/-}, and antibiotic-treated mice, show even stronger such resistance. The question of whether MCAs are important for this resistance may now be investigated by high-fat feeding of MCA-deficient mice. Will these mice respond more like humans?

It is also known that basal bile acid synthesis in mice is about double that seen in humans. Are the FXR antagonistic MCAs important for that species difference? In the present study, there was a clear trend for higher *CYP7A1* mRNA in the WT animals but due to small animal numbers, this did not reach statistical significance. Another issue that also will be of interest is how MCA-deficient mice will respond to bile duct ligation. Will bile acid synthesis be suppressed as in humans or will it be induced as in WT mice? These and many other questions to clarify the physiologic functions of MCAs should now be possible to answer. 

- Russell, D. W. 2003. The enzymes, regulation, and genetics of bile acid synthesis. *Annu. Rev. Biochem.* **72**: 137–174.
- Botham, K. M., and G. S. Boyd. 1983. The metabolism of chenodeoxycholic acid to beta-muricholic acid in rat liver. *Eur. J. Biochem.* **134**: 191–196.
- Takahashi, S., T. Fukami, Y. Masuo, C. N. Brocker, C. Xie, K. W. Krausz, R. Wolf, C. C. Henderson, and F. J. Gonzales. 2016. *Cyp2c70* is responsible for the species difference in bile acid metabolism between mice and humans. *J. Lipid Res.* **57**: 2130–2137.
- Li-Hawkins, J., M. Gafvels, M. Olin, E. G. Lund, U. Andersson, G. Schuster, I. Bjorkhem, D. W. Russell, and G. Eggertsen. 2002. Cholic acid mediates negative feedback regulation of bile acid synthesis in mice. *J. Clin. Invest.* **110**: 1191–1200.
- Hu, X., Y. Bonde, G. Eggertsen, and M. Rudling. 2014. Muricholic bile acids are potent regulators of bile acid synthesis via a positive feedback mechanism. *J. Intern. Med.* **275**: 27–38.
- Zhang, Y., and C. D. Klaassen. 2010. Effects of feeding bile acids and a bile acid sequestrant on hepatic bile acid composition in mice. *J. Lipid Res.* **51**: 3230–3242.
- Makishima, M., A. Y. Okamoto, J. J. Repa, H. Tu, R. M. Learned, A. Luk, M. V. Hull, K. D. Lustig, D. J. Mangelsdorf, and B. Shan. 1999. Identification of a nuclear receptor for bile acids. *Science*. **284**: 1362–1365.
- Parks, D. J., S. G. Blanchard, R. K. Bledsoe, G. Chandra, T. G. Consler, S. A. Kliewer, J. B. Stimmel, T. M. Willson, A. M. Zavacki, D. D. Moore, et al. 1999. Bile acids: natural ligands for an orphan nuclear receptor. *Science*. **284**: 1365–1368.
- Howard, W. R., J. A. Pospisil, E. Njolito, and D. J. Noonan. 2000. Catabolites of cholesterol synthesis pathways and forskolin as activators of the farnesoid X-activated nuclear receptor. *Toxicol. Appl. Pharmacol.* **163**: 195–202.
- Sayin, S. I., A. Wahlstrom, J. Felin, S. Jantti, H. U. Marschall, K. Bamberg, B. Angelin, T. Hyotylainen, M. Oresic, and F. Backhed. 2013. Gut microbiota regulates bile acid metabolism by reducing the levels of tauro-beta-muricholic acid, a naturally occurring FXR antagonist. *Cell Metab.* **17**: 225–235.
- Rabot, S., M. Membrez, A. Bruneau, P. Gerard, T. Harach, M. Moser, F. Raymond, R. Mansourian, and C. J. Chou. 2010. Germ-free C57BL/6J mice are resistant to high-fat-diet-induced insulin resistance and have altered cholesterol metabolism. *FASEB J.* **24**: 4948–4959.
- Bonde, Y., G. Eggertsen, and M. Rudling. 2016. Mice Abundant in Muricholic Bile Acids Show Resistance to Dietary Induced Steatosis, Weight Gain, and to Impaired Glucose Metabolism. *PLoS One*. **11**: e0147772.
- Kaur, A., J. V. Patankar, W. de Haan, P. Ruddle, N. Wijesekara, A. K. Groen, C. B. Verchere, R. R. Singaraja, and M. R. Hayden. 2015. Loss of *Cyp8b1* improves glucose homeostasis by increasing GLP-1. *Diabetes*. **64**: 1168–1179.
- Murphy, C., P. Parini, J. Wang, I. Bjorkhem, G. Eggertsen, and M. Gafvels. 2005. Cholic acid as key regulator of cholesterol synthesis, intestinal absorption and hepatic storage in mice. *Biochim. Biophys. Acta.* **1735**: 167–175.
- Wang, D. Q., S. Tazuma, D. E. Cohen, and M. C. Carey. 2003. Feeding natural hydrophilic bile acids inhibits intestinal cholesterol absorption: studies in the gallstone-susceptible mouse. *Am. J. Physiol. Gastrointest. Liver Physiol.* **285**: G494–G502.
- Dueland, S., J. Reichen, G. T. Everson, and R. A. Davis. 1991. Regulation of cholesterol and bile acid homeostasis in bile-obstructed rats. *Biochem. J.* **280**: 373–377.